

B2.1 Introduction to Representation Theory  
Problem Sheet 4, MT 2017

1. Find the character table of the alternating group  $A_5$ . (It may be helpful to remember that  $A_5$  acts as the group rotations of the regular icosahedron. You may also think about restriction/induction between  $A_5$  and  $S_5$  if it helps.)
2. A conjugacy class  $g^G$  of a group  $G$  is called real if  $g^G = (g^{-1})^G$  i.e., if  $g$  is conjugate to  $g^{-1}$ . A character  $\chi$  of  $G$  is called real if  $\chi(x) \in \mathbb{R}$  for all  $x \in G$ . Prove that the number of real conjugacy classes of a finite group is equal to the number of irreducible real characters. [Hint: Compute the dimension of the complex vector space

$$V := \{f : G \rightarrow \mathbb{C} \mid f(g) = f(h^{-1}gh) = f(g^{-1}) \quad \forall g, h \in G\}$$

in two different ways.

3. Let  $G$  be a finite group with an irreducible representation  $\rho : G \rightarrow GL(2, \mathbb{C})$ .
  - (a) Prove that  $G$  has an element  $a$  of order 2.
  - (b) For  $a$  as above show that either  $\det \rho(a) \neq 1$  or else  $\rho(a)$  is central in  $GL(2, \mathbb{C})$ .
  - (c) Deduce that a finite simple group cannot have an irreducible representation of degree 2.
4. Prove that every finite group  $G$  has a faithful representation. Which finite abelian groups have a faithful *irreducible* representation?
5. Recall from the lectures that an element  $e$  of an algebra  $A$  is called an idempotent if  $e^2 = e$ . Let  $G$  be a finite group and suppose  $V$  is a simple  $\mathbb{C}G$ -module. Define

$$e_V = \frac{\dim V}{|G|} \sum_{g \in G} \overline{\chi_V(g)} g \in \mathbb{C}G.$$

- (a) Prove that  $e_V$  is an element of the centre of  $\mathbb{C}G$ .
  - (b) Let  $V'$  be a simple  $\mathbb{C}G$ -module. Prove that  $e_V$  acts on  $V'$  by 0 if  $V' \not\cong V$  and it acts by the identity on  $V$ .
  - (c) Prove that if  $\{V_i : 1 \leq i \leq n\}$  is the set of irreducible  $G$ -representations (up to isomorphism) and  $e_i = e_{V_i}$ , then  $e_i^2 = e_i$  and  $e_i \cdot e_j = 0$  in  $\mathbb{C}G$ . How does this relate to the Artin-Wedderburn Theorem?
6. Determine the restriction of the standard representation of  $S_4$  to  $S_3$ . Compute the induced of the trivial representation of  $S_3$  to  $S_4$ . Use this to illustrate Frobenius reciprocity.

7. Decompose into irreducible  $G$ -representations the induced representation  $\text{Ind}_H^G W$  where  $G = S_4$  and
- $H = \langle (1234) \rangle$  and  $W = \mathbb{C}v$  is the one-dimensional representation defined by  $(1234) \cdot v = iv$ , where  $i = \sqrt{-1}$ .
  - $H = \langle (123) \rangle$  and  $W = \mathbb{C}v$  is the one-dimensional representation defined by  $(123) \cdot v = e^{2\pi i/3}v$ .
8. (*optional*) Here is another result of Burnside: Let  $V$  be an irreducible representation of a finite group  $G$  and assume that  $\dim V > 1$ . Prove that  $\chi_V$  takes the value 0 on some conjugacy class of  $G$ . (Hint: assume first that  $\chi_V$  takes integer values.)
9. (*optional*) Suppose that  $V$  is a faithful representation of  $G$ . Show that every irreducible representation of  $G$  appears in some tensor power  $V^{\otimes n} = V \otimes V \otimes \cdots \otimes V$  of  $V$ . (Hint: for an arbitrary irreducible character  $\chi$ , consider the infinite series  $\sum_{n \geq 0} \langle \chi, \chi_{V^{\otimes n}} \rangle_G t^n$ , where  $t$  is an indeterminate.)
10. (*optional*) Which irreducible representations of  $S_n$  remain irreducible when restricted to  $A_n$ ? Which irreducible representations of  $S_n$  are induced from  $A_n$ ?